

New perspectives on volcanic clouds from the A-Train

Simon A. Carn<sup>1</sup>, Fred J. Prata<sup>2</sup>, Kai Yang<sup>3</sup>, and Nickolay A. Krotkov<sup>3</sup>

- 1. Michigan Technological University, Houghton, MI, USA
- 2. Norwegian Institute for Air Research, Kjeller, Norway
- 3. GEST Center, UMBC, Baltimore, MD, USA



## Volcanic clouds: 'dirty thunderstoms'

#### Volcanic eruption cloud


- Vertically extensive
- Short-lived (hours)
- Overshooting top
- Stratospheric penetration
- Lightning
- Umbrella cloud (gravity current)
- Overseeded' with ice nuclei (ash)
- Ash (µm cm), hydrometeors
- Layered structure
- 'Downdraft' during ashfall?

#### Single-cell Cb


- Vertically extensive
- Short-lived (hours)
- Overshooting top
- Rise to tropopause
- Anvil: ice-rich blow-off
- Hydrometeors (rain hail)
- Lightning
- Downdrafts








#### Aviation hazards from volcanic clouds





- Acute hazards
  - Engine failure due to melted ash
  - Abrasion of windshield
  - Effects on avionics
- Secondary hazards
  - Corrosion by ash, sulfuric acid
- Mitigation
  - Immediate detection of fresh volcanic clouds
  - Tracking/forecast of cloud position and <u>altitude</u>



From: Volcanoes; Crucibles of Change, Princeton U. Press, Princeton, 1997.

#### Two near-disastrous volcanic cloud encounters





"Ladies and gentlemen, this is your captain speaking. We have a small problem. All four engines have stopped. We are doing our damnedest to get them under control. I trust you are not in too much distress." Capt. Eric Moody, BA Flight 9

BA 009 (B747), Kuala Lumpur – Perth, 24 June 1982

KLM 867 (B747), Amsterdam – Tokyo, 15 December 1989 (Redoubt, AK)


~2 g/m<sup>3</sup> ash?

## Volcanic ash sedimentation




Mammatus lobes on the Mount St. Helens volcanic cloud over Ephrata, WA, 18 May 1980 (© D. Miller)


#### Hydrometeor-enhanced sedimentation of volcanic ash



- Ash aggregation prematurely removes fine ash that would be an aviation hazard
- How do we explain secondary ash thickness maxima and ash aggregate fallout?
- Can CloudSat W-band radar detect the bright band?


# Exploiting A-Train synergy for volcanic cloud studies









Mass: 195.725 kt; Area: 1400753 km2; SO2 max: 32.71 DU at lon: -74.40 lat: 11.42






Mass: 188.536 kt; Area: 1880299 km2; SO2 max: 23.66 DU at lon: -81.33 lat: 14.47






Mass: 171.187 kt; Area: 2410030 km2; SO2 max: 19.89 DU at lon: -93.38 lat: 13.66






Mass: 159.341 kt; Area: 2532097 km2; SO2 max: 20.18 DU at lon: -107.02 lat: 12.14







Mass: 168.052 kt; Area: 3126612 km2; SO2 max: 18.41 DU at lon: -114.18 lat: 13.47







Mass: 146.936 kt; Area: 3468046 km2; SO2 max: 12.14 DU at lon: -121.13 lat: 13.78






















Mass: 98.493 kt; Area: 1701411 km²; SO<sub>2</sub> max: 6.22 DU at lon: -145.52 lat: 12.46








Mass: 107.076 kt; Area: 2034725 km2; SO2 max: 6.74 DU at lon: -146.34 lat: 13.83














[Carn et al., 2007; Prata et al., 2007]






Mass: 79.103 kt; Area: 2352265 km2; SO2 max: 3.01 DU at lon: 155.20 lat: 11.69


[Carn et al., 2007; Prata et al., 2007]







Mass: 71.854 kt; Area: 2170677 km2; SO2 max: 2.94 DU at lon: 151.98 lat: 8.57














Mass: 45.728 kt; Area: 1734106 km²; SO<sub>2</sub> max: 2.12 DU at lon: 126.13 lat: 4.70




## CALIPSO first light – 7 June 2006



#### CALIPSO – OMI – MLS: 8 June 2006




## CALIPSO profiles through SHV aerosol cloud




- Aerosol layer non-depolarizing
- Liquid sulfate aerosol dominant


## The 2010 eruption of Eyjafjallajökull (Iceland)



## Volcanic Ash Advisory (VAA) and OMI AI – April 16



### A-Train measurements – April 16



## Eyjafjallajökull: 'strong' and 'weak' eruption plumes






 $H = 1.67Q^{0.259}$ 

- Empirical relationship of mass eruption rate
   (Q) to plume height (H) for 'strong' plumes
- Assumes a single injection height
- Does not account for environmental factors (e.g., wind, humidity), gravity current intrusions etc.




- When does advection-diffusion take over?
- -> Initialize models with accurate vertical profile information

## A-Train observations of Eyjafjallajökull plume – May 8, 2010




## Eyjafjallajökull eruption plume – April 15, 2010




• NB: OMI row anomaly coincides with CALIPSO/CloudSat track

#### CALIPSO + CloudSat + OMI: April 15, 2010



## A-Train observations: Kasatochi, August 9, 2008




- Aqua MODIS
- Largest volcanic SO<sub>2</sub> emission since 1991 (~2 Tg)
- Negligible climate impact


#### JGR Special issue

http://www.agu.org/journals/jd/special\_sections.shtml?collectionCode=VOLCANOES1&journalCode=JD

## A-Train observations: Kasatochi, August 9, 2008




## Forward trajectories for the Kasatochi volcanic cloud



## A-Train observations: Chaitén (Chile), May 2008









- Eruption at ~0800
   local time on 6 May
- High wind shear environment




### IR ash composition retrievals – the 'Concavity' signature




#### Volcanic ash tracking with the AIRS concavity algorithm




### A-Train observations: Chaitén (Chile), May 7, 2008



## Monitoring tropospheric volcanic plumes with CALIPSO





# Sarychev Peak (Kurile Is) eruption




A-Train synergy compensates for data gap and improves SO<sub>2</sub> loading measurements

## Hypothetical A-Train eruption monitoring system







# Summary

- The A-Train is providing unprecedented observations of the vertical distribution of trace gases, aerosols and hydrometeors in volcanic clouds
- New insights into ashfall processes and stratospheric aerosol evolution in volcanic clouds are anticipated (see talk by Mike Fromm)
- Measurements of ash cloud altitude and thickness (e.g., by CALIPSO)
  are of prime importance for aviation hazard mitigation
- Currently, data latency of most A-Train products (and spatial coverage of the active sensors) is the main impediment to operational application for aviation hazard mitigation
- The A-Train greatly facilitates validation/intercomparison of satellite data products (e.g., SO<sub>2</sub> column measurements)

#### Acknowledgments:







